

Anti-Diabetic Research on Bitter Melon (*Momordica charantia*)

- Niu, Y., et al. "Comprehensive studies on the regulation of type 2 diabetes by cucurbitane-type triterpenoids in *Momordica charantia* L.: Insights from network pharmacology and molecular docking and dynamics." *Pharmaceuticals* (Basel). 2025 Mar; 18(4): 474.
- Yadav, R., et al. "Exploring the therapeutic potential of *Momordica charantia* in targeting protein kinase C delta (PRKCD) for type 2 diabetes mellitus: Insights from network pharmacology, molecular docking, and molecular dynamics simulations." *In Silico Pharmacol.* 2025 Jul; 13(2): 99.
- Mes, J., et al. "Bitter gourd (*Momordica charantia* L.) supplementation for twelve weeks improves biomarkers of glucose homeostasis in a prediabetic population." *J. Ethnopharmacol.* 2025 May; 347: 119756.
- Okechukwu, U., et al. "Harnessing plant metabolic pathways for innovative diabetes management: unlocking the therapeutic potential of medicinal plants." *Plant Signal Behav.* 2025 Dec; 20(1): 2486076.
- Romdhoni, M., et al. "Antidiabetic activity of *Momordica charantia* extracts through incretin pathway in streptozotocin-nicotinamide induced diabetic rat depends on dose differences." *Avicenna J. Med. Biotechnol.* 2025 Jan-Mar; 17(1): 47-55.
- Gautam, V., et al. "Exploring the therapeutic potential of three cucurbit plants involving *in vivo* diabetes screening." *Cureus.* 2025 Feb; 17(2): e78861.
- Zhang, Y., et al. Polysaccharide from *Momordica charantia* L. alleviates Type 2 Diabetes mellitus in mice by activating the IRS1/PI3K/Akt and AMPK signaling pathways and regulating the gut microbiota." *J. Agric. Food Chem.* 2025 Mar; 73(12): 7298-7309.
- Hsiang, C., et al. "Synergistic action of a *Momordica charantia*-abundant peptide with insulin regulates blood glucose metabolism in db/db mice via a novel second-hit mechanism on insulin receptor activation." *Int. J. Biol. Macromol.* 2025 Jun; 311 (Pt 2): 143531.
- Niu, Y., et al. "Comprehensive studies on the regulation of type 2 diabetes by cucurbitane-type triterpenoids in *Momordica charantia* L.: Insights from network pharmacology and molecular docking and dynamics. *Pharmaceuticals* (Basel). 2025 Mar; 18(4): 474.
- Zhang, Y., et al. "Ultrasound-assisted extraction, purification, structural characterization, and hypoglycemic activities of a polysaccharide from *Momordica charantia* L." *Int. J. Biol. Macromol.* 2025 Jan; 285: 138300.
- Famuyiwa, S., et al. "Investigating the multitargeted anti-diabetic potential of cucurbitane-type triterpenoid from *Momordica charantia*: an LC-MS, docking-based MM\GBSA and MD simulation study." *J. Biomol. Struct. Dyn.* 2025 Feb; 43(3): 1159-1170.

Hung, W., et al. Exploring the antidiabetic and antihypertensive potential of peptides derived from bitter melon seed hydrolysate." *Biomedicines*. 2024 Oct; 12(11): 2452.

Zhao, D., et al. "Metabolomics revealed the effects of *Momordica charantia* L. saponins on diabetic hyperglycemia and wound healing in mice." *Foods*. 2024 Oct; 13(19): 3163.

Rivera, R., et al. "Applying network pharmacology and molecular docking in the screening for molecular mechanisms of ampalaya (*Momordica charantia* L.) and banaba (*Lagerstroemia speciosa* L.) against Type 2 Diabetes mellitus." *Acta Med. Philipp.* 2024 May; 58(8): 108-124.

Fan, Z., et al. "Integration of microRNA and mRNA analyses depicts the potential roles of *Momordica charantia* saponin administration in insulin resistance of juvenile common carp (*Cyprinus carpio*) fed with a high-starch diet." *Front. Mol. Biosci.* 2023 Apr; 10: 1054949.

Sun, K., et al. "Effects of dietary wild bitter melon (*Momordica charantia* var. abbreviate Ser.) extract on glucose and lipid metabolism in HFD/STZ-induced type 2 diabetic rats." *J. Ethnopharmacol.* 2023 Apr; 306: 116154.

Ali, A., et al. "Effect of *Momordica charantia* on insulin immune-reactive pancreatic beta cells and blood glucose levels in streptozotocin-induced diabetic rats." *J. Nutr. Sci. Vitaminol.* (Tokyo). 2022; 68(5): 438-445.

Hussain, F., et al. "In vitro and in vivo study of inhibitory potentials of α -glucosidase and acetylcholinesterase and biochemical profiling of *M. charantia* in alloxan-induced diabetic rat models." *Am. J. Transl. Res.* 2022 Jun; 14(6): 3824-3839.

Tehseen, I., et al. "Antidiabetic and hepato-renal protective effects of medicinal plants in STZ induced diabetic rats." *Braz. J. Biol.* 2022 Jun; 84: e260189.

Xu, B., et al. "Bioactives of *Momordica charantia* as potential anti-diabetic/hypoglycemic agents." *Molecules*. 2022 Mar; 27(7): 2175.

Sajadimajd, S., et al. "Modulation of notch signaling and angiogenesis via an isolated polysaccharide from *Momordica charantia* in diabetic rats." *J. Food. Biochem.* 2022 Feb; 46(2): e14033.

Makena, W., et al. "Genistein and *Momordica charantia* L. prevent oxidative stress and upregulate proglucagon and insulin receptor mRNA in diabetic rats." *Appl. Physiol. Nutr. Metab.* 2022 Jan; 47(1): 41-50.

Arif, R., et al. "Molecular docking and simulation studies of antidiabetic agents devised from hypoglycemic Polypeptide-P of *Momordica charantia*." *Biomed. Res. Int.* 2021 Sep; 2021: 5561129.

Liu, Y., et al. "Saponins of *Momordica charantia* increase insulin secretion in INS-1 pancreatic β -cells via the PI3K/Akt/FoxO1 signaling pathway." *Endocrinol. Diabetes Nutr.* (Engl Ed). 2021 May; 68(5): 329-337.

Mahwish, F., et al. "Bitter Melon (*Momordica charantia* L.) fruit bioactives charantin and vicine potential for diabetes prophylaxis and treatment." *Plants* (Basel). 2021 Apr; 10(4): 730.

Kulkarni, P., et al. "Bioanalytical method development for momordicin and its application to long-term pharmacokinetics in diabetic rats." *Drug Dev. Ind. Pharm.* 2021 Jul; 47(7): 1064-1071.

Mahrosh, H., et al. "Investigation of hypoglycemic peptides derived from conserved regions of adMc1 to reveal their antidiabetic activities." *Biomed. Res. Int.* 2021 Mar; 2021: 5550180.

Kulkarni, P., et al. "Isolation, characterisation and investigation of *in vitro* antidiabetic and antioxidant activity of phytoconstituents from fruit of *Momordica charantia* Linn." *Nat. Prod. Res.* 2021 Mar; 35(6): 1035-1037.

Shimada, T., et al. "Bitter melon fruit extract enhances intracellular ATP production and insulin secretion from rat pancreatic β -cells." *Br. J. Nutr.* 2022 Feb; 127(3): 377-383.

Chang, C., et al. "Bitter melon extract yields multiple effects on intestinal epithelial cells and likely contributes to anti-diabetic functions." *Int. J. Med. Sci.* 2021 Feb; 18(8): 1848-1856.

Perera, W., et al. "Anti-inflammatory, antidiabetic properties and *in silico* modeling of cucurbitane-type triterpene glycosides from fruits of an Indian cultivar of *Momordica charantia* L. molecules." 2021 Feb; 26(4): 1038.

Lee, Y., et al. "Metabolite profile of cucurbitane-type triterpenoids of bitter melon (fruit of *Momordica charantia*) and their inhibitory activity against protein tyrosine phosphatases relevant to insulin resistance." *J. Agric. Food Chem.* 2021 Feb; 69(6): 1816-1830.

Makena, W., et al. "*Momordica charantia* L. fruit and Genistein ameliorates type 2 diabetes in rats by preventing lipid accumulation, insulin resistance and enhancing beta cell function." *J. Diabetes Metab. Disord.* 2020 Oct; 19(2): 1303-1310.

Gao, Y., et al. "Cucurbitane-type triterpene glycosides from *Momordica charantia* and their α -glucosidase inhibitory activities." *Nat. Prod. Bioprospect.* 2020 Jun; 10(3): 153-161.

Dwijayanti, D. et al. "Bitter melon fruit extract has a hypoglycemic effect and reduces hepatic lipid accumulation in ob/ob mice." *Phytother. Res.* 2020 Jun; 34(6): 1338-1346.

Shivanagoudra, S., et al. "*In vitro* and *in silico* elucidation of antidiabetic and anti-inflammatory activities of bioactive compounds from *Momordica charantia* L." *Bioorg. Med. Chem.* 2019 Jul; 27(14): 3097-3109.

Wang, Q., et al. "Comparison of antidiabetic effects of saponins and polysaccharides from *Momordica charantia* L. in STZ-induced type 2 diabetic mice." *Biomed. Pharmacother.* 2019 Jan; 109: 744-750.

Alam, F., et al. "Enzymes inhibitors from natural sources with antidiabetic activity: A review." *Phytother. Res.* 2019 Jan; 33(1): 41-54.

Zhang, C., et al. "Preparation of a *Momordica charantia* L. polysaccharide chromium (III) complex and its anti-hyperglycemic activity in mice with streptozotocin-induced diabetes." *Int. J. Bio. Macromol.* 2019 Feb; 122: 619-627.

Malekshahi, H., et al. "*Momordica charantia* reverses type II diabetes in rat." *J. Food. Biochem.* 2019 Nov; 43(11): e13021.

Peter, E., et al. "*Momordica charantia* L. lowers elevated glycaemia in Type 2 Diabetes mellitus

patients: Systematic review and meta-analysis." *J. Ethnopharmacol.* 2019 Mar; 231: 311-324.

Farah, N., et al. "Phenolic acid profiling and antiglycation studies of leaf and fruit extracts of tyrosine primed *Momordica charantia* seeds for possible treatment of diabetes mellitus." *Pak. J. Pharm. Sci.* 2018 Nov; 31(6 (Supplementary): 2667-2672.

Chanda, J., et al. "Study of pancreatic lipase inhibition kinetics and LC-QTOF-MS-based identification of bioactive constituents of *Momordica charantia* fruits." *Biomed. Chromatogr.* 2018 Dec 9: e4463.

Peter, E., et al. "Safety and efficacy of *Momordica charantia* Linnaeus in pre-diabetes and type 2 diabetes mellitus patients: A systematic review and meta-analysis protocol." *Syst. Rev.* 2018 Nov; 7(1):192.

Fuller, S., et al. "Potential adverse effects of botanical supplementation in high-fat-fed female mice." *Biol. Sex. Differ.* 2018 Sep 12; 9(1): 41.

Bai, W., et al. "Transcriptomic analysis of *Momordica charantia* polysaccharide on streptozotocin-induced diabetic rats." *Gene.* 2018 Oct; 675: 208-216.

Hussain, F., et al. "Study of selected therapeutic potentials of *Momordica charantia* fruit." *J. Biol. Regul. Homeost. Agents.* 2018 Jul-Aug; 32(4): 859-862.

Bhat, G., et al. "Correction to: GLP-I secretion in healthy and diabetic Wistar rats in response to aqueous extract of *Momordica charantia*." *BMC Complement. Altern. Med.* 2018 Jun 12; 18(1): 82.

Bai, J., et al. "Obese rats supplemented with bitter melon display marked shifts in the expression of genes controlling inflammatory response and lipid metabolism by RNA-Seq analysis of colonic mucosa." *Genes Genomics.* 2018 Jun; 40(6): 561-567.

Bhat, G., et al. "Correction to: GLP-I secretion in healthy and diabetic Wistar rats in response to aqueous extract of *Momordica charantia*." *BMC Complement. Altern. Med.* 2018 Jun; 18(1):175.

Bhat, G., et al. "GLP-I secretion in healthy and diabetic Wistar rats in response to aqueous extract of *Momordica charantia*." *BMC Complement. Altern. Med.* 2018 May; 18(1): 162.

Elekofehintj, O., et al. "Potential use of bitter melon (*Momordica charantia*) derived compounds as antidiabetics: In silico and *in vivo* studies." *Pathophysiology.* 2018 Dec; 25(4): 327-333.

Deshaware, S., et al. "Debittering of bitter gourd juice using β -cyclodextrin: Mechanism and effect on antidiabetic potential." *Food Chem.* 2018 Oct; 262: 78-85.

Zhang, C., et al. "Characterization of *Momordica charantia* L. polysaccharide and its protective effect on pancreatic cells injury in STZ-induced diabetic mice." *Int. J. Biol. Macromol.* 2018 Aug; 115: 45-52.

Sandikapura, M., et al. "Comparative antioxidant and antidiabetic effects of *Syzygium polyanthum* leaf and *Momordica charantia* fruit extracts." *Pak. J. Pharm. Sci.* 2018 Mar; 31(2(Suppl.)): 623-635.

Han, J., et al. "Cucurbitane triterpenoids from the fruits of *Momordica charantia* improve insulin sensitivity and glucose homeostasis in streptozotocin-induced diabetic mice." *Mol. Nutr. Food*

Res. 2018 Apr; 62(7): e1700769.

Krawinkel, M., et al. "Bitter gourd reduces elevated fasting plasma glucose levels in an intervention study among prediabetics in Tanzania." *J. Ethnopharmacol.* 2018 Apr; 216: 1-7.

Fachinan, R., et al. "Effectiveness of antihyperglycemic effect of *Momordica charantia*: implication of T-cell cytokines." *Evid. Based Complement. Alternat. Med.* 2017; 2017: 3707046.

Gao, Z., et al. "New insights into the mechanisms of Chinese herbal products on diabetes: A focus on the "bacteria-mucosal immunity-inflammation-diabetes axis." *J. Immunol. Res.* 2017; 2017: 1813086.

Ma, C., et al. "*Momordica charantia* extracts ameliorate insulin resistance by regulating the expression of SOCS-3 and JNK in type 2 diabetes mellitus rats." *Pharm. Biol.* 2017 Dec; 55(1): 2170-2177.

Ramachandra Shobha, C., et al. "Fifty percent ethanolic extract of *Momordica charantia* inhibits adipogenesis and promotes adipolysis in 3T3-L1 pre-adipocyte cells." *Rep. Biochem. Mol. Biol.* 2017 Oct; 6(1): 22-32.

Chen, G., et al. "The anti-adiposity effect of bitter melon seed oil is solely attributed to its fatty acid components." *Lipids Health Dis.* 2017 Sep; 16(1): 186.

Lo, H., et al. "Gastro-resistant insulin receptor-binding peptide from *Momordica charantia* improved the glucose tolerance in streptozotocin-induced diabetic mice via insulin receptor signaling pathway." *J. Agric. Food Chem.* 2017 Oct; 65(42): 9266-9274.

Chong, W., et al. "*Momordica charantia*: A new strategic vision to improve the therapy of endoplasmic reticulum stress." *Curr. Pharm. Des.* 2017; 23(16): 2333-2343.

Mahmoud, M., et al. "Studies on the antidiabetic activities of *Momordica charantia* fruit juice in streptozotocin-induced diabetic rats." *Pharm. Biol.* 2017 Dec; 55(1): 758-765.

Jiang, B., et al. "Antidiabetic activities of a cucurbitane type triterpenoid compound from *Momordica charantia* in alloxan induced diabetic mice." *Mol. Med. Rep.* 2016 Nov; 14(5): 4865-4872.

Güdr, A., et al. "Influence of total anthocyanins from bitter melon (*Momordica charantia* Linn.) as antidiabetic and radical scavenging agents." *Iran J. Pharm. Res.* 2016 Winter; 15(1): 301-9.

Boone, C., et al. "Acute effects of a beverage containing bitter melon extract (CARELA) on postprandial glycemia among prediabetic adults." *Nutr. Diabetes.* 2017 Jan; 7(1): e241.

Aljohi, A., et al. "Antiglycation and antioxidant properties of *Momordica charantia*." *PLoS One.* 2016 Aug; 11(8): e0159985.

Zhu, Y., et al. "Effects of bitter melon (*Momordica charantia* L.) on the gut microbiota in high fat diet and low dose streptozotocin-induced rats." *Int. J. Food Sci. Nutr.* 2016 Sep; 67(6): 686-95.

Chandru, S., et al. "Evaluation of protein kinase C β and PPAR γ activity in diabetic rats supplemented with *Momordica charantia*." *J. Clin. Diagn. Res.* 2016 Apr; 10(4): BF01-4.

Lo, H., et al. "Identification of the bioactive and consensus peptide motif from *Momordica charantia* insulin receptor-binding protein." *Food Chem.* 2016 Aug; 204: 298-305.

Tan, H., et al. "Polysaccharide with antioxidant, α -amylase inhibitory and ACE inhibitory activities from *Momordica charantia*." *Int. J. Biol. Macromol.* 2016 Apr; 85: 487-96.

Burnett, A., et al. "Investigation of the blood glucose lowering potential of the Jamaican *Momordica charantia* (Cerasee) fruit in Sprague-Dawley rats." *West Indian Med. J.* 2015 Sep; 64(4): 315-9.

Rudá-Kučerová, J., et al. "[Effectiveness of phytotherapy in supportive treatment of type 2 Diabetes mellitus III. *Momordica* (*Momordica charantia*)]." *Ceska Slov. Farm.* 2015 Sep; 64(4): 126-32.

Xu, X., et al. "Anti-diabetic properties of *Momordica charantia* L. polysaccharide in alloxan-induced diabetic mice." *Int. J. Biol. Macromol.* 2015 Nov; 81: 538-43.

Yang, B., et al. "Design, synthesis and biological evaluation of novel peptide MC2 analogues from *Momordica charantia* as potential anti-diabetic agents." *Org. Biomol. Chem.* 2015 Apr; 13(15): 4551-61.

Chen, J., et al. "The antigluconeogenic activity of cucurbitacins from *Momordica charantia*." *Planta Med.* 2015 Mar; 81(4): 327-32.

Yang, S., et al. "Preventive effects of bitter melon (*Momordica charantia*) against insulin resistance and diabetes are associated with the inhibition of NF- κ B and JNK pathways in high-fat-fed OLETF rats." *J. Nutr. Biochem.* 2015 Mar; 26(3): 234-40.

Tahira, S., et al. "Antidiabetic evaluation of *Momordica charantia* L fruit extracts." *West Indian Med. J.* 2014 Aug; 63(4): 294-9.

Medagama, A., et al. "The use of complementary and alternative medicines (CAMs) in the treatment of diabetes mellitus: Is continued use safe and effective?" *Nutr. J.* 2014 Oct; 13: 102.

Bachok, M., et al. "Effectiveness of traditional Malaysian vegetables (ulam) in modulating blood glucose levels." *Asia Pac. J. Clin. Nutr.* 2014; 23(3): 369-76.

Lo, H., et al. "A novel insulin receptor-binding protein from *Momordica charantia* enhances glucose uptake and glucose clearance *in vitro* and *in vivo* through triggering insulin receptor signaling pathway." *J. Agric. Food Chem.* 2014 Sep; 62(36): 8952-61.

Wu, S., et al. "Transport in Caco-2 cell monolayers of antidiabetic cucurbitane triterpenoids from *Momordica charantia* fruits." *Planta Med.* 2014 Jul; 80(11):907-11.

Tian, M., et al. "Molecular diversity and hypoglycemic polypeptide-P content of *Momordica charantia* in different accessions and different seasons." *J. Sci. Food Agric.* 2015 Apr; 95(6): 1328-35.

Nkambo, W., et al. "*In vivo* hypoglycemic effect of methanolic fruit extract of *Momordica charantia* L." *Afr. Health Sci.* 2013 Dec; 13(4): 933-9.

Wang, H., et al. "Differential anti-diabetic effects and mechanism of action of charantin-rich extract of Taiwanese *Momordica charantia* between type 1 and type 2 diabetic mice." *Food Chem. Toxicol.* 2014 Jul; 69: 347-56.

França, E., et al. "Effects of *Momordica charantia* L. on the blood rheological properties in

diabetic patients." *Biomed. Res. Int.* 2014; 2014: 840379.

Efird, J., et al. "Potential for improved glycemic control with dietary *Momordica charantia* in patients with insulin resistance and pre-diabetes." *Int. J. Environ. Res. Public Health.* 2014 Feb; 11(2): 2328-45.

Huang, T., et al. "Role of GLP-1 in the hypoglycemic effects of wild bitter gourd." *Evid. Based Complement. Alternat. Med.* 2013; 2013: 625892.

Yu, Y., et al. "Bitter melon extract attenuating hepatic steatosis may be mediated by FGF21 and AMPK/Sirt1 signaling in mice." *Sci. Rep.* 2013 Nov; 3: 3142.

Patel, D., et al. "An overview on antidiabetic medicinal plants having insulin mimetic property." *Asian Pac J Trop Biomed.* 2012 Apr; 2(4):320-30.

Hsiao, P., et al. "Antiproliferative and hypoglycemic cucurbitane-type glycosides from the fruits of *Momordica charantia*." *J. Agric. Food Chem.* 2013 Mar; 61(12): 2979-86.

Lo, H., et al. "*Momordica charantia* and its novel polypeptide regulate glucose homeostasis in mice via binding to insulin receptor." *J. Agric. Food Chem.* 2013 Mar; 61(10): 2461-8.

Hu XJ, et al. "α-Glucosidase inhibitors via green pathway: biotransformation for bicoumarins catalyzed by *Momordica charantia* peroxidase." *J. Agric. Food Chem.* 2013 Feb; 61(7): 1501-8.

Choudhary, S., et al. "Comprehensive evaluation of anti-hyperglycemic activity of fractionated *Momordica charantia* seed extract in alloxan-induced diabetic rats." *Evid. Based. Complement. Alternat. Med.* 2012; 2012: 293650.

Ooi, C., et al. "*Momordica charantia* for type 2 diabetes mellitus." *Cochrane Database Syst. Rev.* 2012 Aug; 8: CD007845.

Ahmad, Z., et al. "*In vitro* anti-diabetic activities and chemical analysis of polypeptide-k and oil isolated from seeds of *Momordica charantia* (bitter gourd)." *Molecules.* 2012 Aug; 17(8): 9631-40.

Mahomoodally, M., et al. "Traditional medicinal herbs and food plants have the potential to inhibit key carbohydrate hydrolyzing enzymes *in vitro* and reduce postprandial blood glucose peaks *in vivo*." *Scientific World Journal.* 2012; 2012: 285284.

Deng, R. "A review of the hypoglycemic effects of five commonly used herbal food supplements." *Recent Pat. Food Nutr. Agric.* 2012 Apr; 4(1): 50-60.

Tsai, C., et al. "Wild bitter gourd improves metabolic syndrome: a preliminary dietary supplementation trial." *Nutr. J.* 2012 Jan; 11: 4.

Chaturvedi, P. "Antidiabetic potentials of *Momordica charantia*: multiple mechanisms behind the effects." *J. Med. Food.* 2012 Feb; 15(2): 101-7

Saha, S., et al. "Antioxidant and anti-inflammatory effect of conjugated linolenic acid isomers against streptozotocin-induced diabetes." *Br. J Nutr.* 2012 Sep; 108(6): 974-83.

Keller, A., et al. "Saponins from the traditional medicinal plant *Momordica charantia* stimulate insulin secretion *in vitro*." *Phytomedicine.* 2011 Dec; 19(1): 32-7.

Akhtar, N., et al. "Pharmaceutical and biopharmaceutical evaluation of extracts from different

plant parts of indigenous origin for their hypoglycemic responses in rabbits." *Acta Pol. Pharm.* 2011 Nov-Dec; 68(6): 919-25.

Blum, A., et al. "*Momordica charantia* extract, a herbal remedy for type 2 diabetes, contains a specific 11 β -hydroxysteroid dehydrogenase type 1 inhibitor." *J. Steroid. Biochem. Mol. Biol.* 2012 Jan; 128(1-2): 51-5.

Singh, J., et al. "Medicinal chemistry of the anti-diabetic effects of *Momordica charantia*: active constituents and modes of actions." *Open Med. Chem. J.* 2011; 5(Suppl 2): 70-7.

Popovich, D., et al. "*Momordica charantia* seed extract reduces pre-adipocyte viability, affects lactate dehydrogenase release, and lipid accumulation in 3T3-L1 cells." *J. Med. Food.* 2011 Mar; 14(3): 201-8.

Feng, E., et al. "Bitter gourd (*Momordica charantia*) is a cornucopia of health: a review of its credited antidiabetic, anti-HIV, and antitumor properties." *Curr. Mol. Med.* 2011 Jul; 11(5): 417-36.

Hafizur, R., et al. "Modulation of pancreatic B-cells in neonatally streptozotocin-induced type 2 diabetic rats by the ethanolic extract of *Momordica charantia* fruit pulp." *Nat Prod Res.* 2011 Feb; 25(4): 353-67.

Wang, Z., et al. "Bioactives from bitter melon enhance insulin signaling and modulate acyl carnitine content in skeletal muscle in high-fat diet-fed mice." *J. Nutr. Biochem.* 2011 Nov; 22(11): 1064-73.

Fuangchan, A., et al. "Hypoglycemic effect of bitter melon compared with metformin in newly diagnosed type 2 diabetes patients." *J. Ethnopharmacol.* 2011 Mar; 134(2): 422-8.

Xie, W., et al. "Diabetes is an inflammatory disease: evidence from traditional Chinese medicines." *Diabetes Obes. Metab.* 2011 Apr; 13(4): 289-301.

Bhujbal, S., et al. "A novel herbal formulation in the management of diabetes." *Int. J. Pharm. Investig.* 2011 Oct; 1(4): 222-6.

Abdollahi, M., et al. "Effects of *Momordica charantia* on pancreatic histopathological changes associated with streptozotocin-induced diabetes in neonatal rats." *Histol. Histopathol.* 2011 Jan; 26(1): 13-21.

Gadang, V., et al. "Dietary bitter melon seed increases peroxisome proliferator-activated receptor- γ gene expression in adipose tissue, down-regulates the nuclear factor- κ B expression, and alleviates the symptoms associated with metabolic syndrome." *J. Med. Food.* 2011 Jan-Feb; 14(1-2): 86-93.

Malik, Z., et al. "Neuroprotective effect of *Momordica charantia* in global cerebral ischemia and reperfusion induced neuronal damage in diabetic mice." *J. Ethnopharmacol.* 2011 Jan; 133(2): 729-34.

Tripathi, U., et al. "Anti-hyperglycemic and anti-oxidative effect of aqueous extract of *Momordica charantia* pulp and *Trigonella foenum graecum* seed in alloxan-induced diabetic rats." *Indian J. Biochem. Biophys.* 2010 Aug; 47(4): 227-33.

Teoh, S., et al. "Histological changes in the kidneys of experimental diabetic rats fed with *Momordica charantia* (bitter gourd) extract." *Rom. J. Morphol. Embryol.* 2010;51(1):91-5.

Babish, J., et al. "Antidiabetic screening of commercial botanical products in 3T3-L1 adipocytes and db/db mice." *J. Med. Food.* 2010 Jun;13(3):535-47.

Xia, X., et al. "Targeting metabolic syndrome: candidate natural agents." *J. Diabetes.* 2010 Dec; 2(4): 243-9.

Klomann, S., et al. "Antidiabetic effects of bitter gourd extracts in insulin-resistant db/db mice." *Br. J. Nutr.* 2010 Dec; 104(11): 1613-20.

Chaturvedi, P., et al. "*Momordica charantia* maintains normal glucose levels and lipid profiles and prevents oxidative stress in diabetic rats subjected to chronic sucrose load." *J. Med. Food.* 2010 Jun; 13(3): 520-7.

Nerurkar, P., et al. "*Momordica charantia* (bitter melon) inhibits primary human adipocyte differentiation by modulating adipogenic genes." *BMC Complement. Altern. Med.* 2010 Jun; 10: 34.

Leung, L., et al. "Anti-diabetic and hypoglycaemic effects of *Momordica charantia* (bitter melon): A mini review." *Br. J. Nutr.* 2009; 102(12): 1703-8.

Tripathi, U., et al. "The plant extracts of *Momordica charantia* and *Trigonella foenum-graecum* have anti-oxidant and anti-hyperglycemic properties for cardiac tissue during diabetes mellitus." *Oxid. Med. Cell Longev.* 2009 Nov-Dec; 2(5): 290-6.

Yadav, M., et al. "Complementary and comparative study on hypoglycemic and antihyperglycemic activity of various extracts of *Eugenia jambolana* seed, *Momordica charantia* fruits, *Gymnema sylvestre*, and *Trigonella foenum graecum* seeds in rats." *Appl. Biochem. Biotechnol.* 2010 Apr; 160(8): 2388-400.

Teoh, S., et al. "A histological study of the structural changes in the liver of streptozotocin-induced diabetic rats treated with or without *Momordica charantia* (bitter gourd)." *Clin Ter.* 2009 Jul-Aug; 160(4): 283-6.

Nahas, R., et al. "Complementary and alternative medicine for the treatment of type 2 diabetes." *Can. Fam. Physician.* 2009; 55(6): 591-6.

Shih, C., et al. "*Momordica charantia* extract on insulin resistance and the skeletal muscle GLUT4 protein in fructose-fed rats." *J. Ethnopharmacol.* 2009 May; 123(1) :82-90.

Inayat-ur-Rahman, M., et al. "Serum sialic acid changes in non-insulin-dependant diabetes mellitus (NIDDM) patients following bitter melon (*Momordica charantia*) and rosiglitazone (Avandia) treatment." *Phytomedicine.* 2009 May; 16(5): 401-5.

Nivitabishekam, S., et al. "Pharmacodynamic interaction of *Momordica charantia* with rosiglitazone in rats." *Chem. Biol. Interact.* 2009 Feb; 177(3): 247-53.

Han, C. "Hypoglycaemic activity of saponin fraction extracted from *Momordica charantia* in PEG/salt aqueous two-phase systems." *Nat. Prod. Res.* 2008; 22(13): 1112-9.

Cheng, H., et al. "A cell-based screening identifies compounds from the stem of *Momordica*

charantia that overcome insulin resistance and activate AMP-activated protein kinase." *J. Agric. Food Chem.* 2008 Aug; 56(16): 6835-43.

Shih, C., et al. "Effects of *Momordica charantia* on insulin resistance and visceral obesity in mice on high-fat diet." *Diabetes Res. Clin. Pract.* 2008 Aug; 81(2): 134-43.

Tan, M., et al. "Antidiabetic activities of triterpenoids isolated from bitter melon associated with activation of the AMPK pathway." *Chem. Biol.* 2008; 15(3): 263-73.

Singh, N., et al. "Regeneration of beta cells in islets of Langerhans of pancreas of alloxan diabetic rats by acetone extract of *Momordica charantia* (Linn.) (bitter gourd) fruits." *Indian J. Exp. Biol.* 2007; 45(12): 1055-62.

Uebanso, T., et al. "Extracts of *Momordica charantia* suppress postprandial hyperglycemia in rats." *J. Nutr. Sci. Vitaminol.* (Tokyo). 2007 Dec; 53(6): 482-8.

Kumar, G., et al. "Modulatory effect of bitter gourd (*Momordica charantia* LINN.) on alterations in kidney heparan sulfate in streptozotocin-induced diabetic rats." *J. Ethnopharmacol.* 2008 Jan 17; 115(2): 276-83.

Sridhar, M., et al. "Bitter gourd (*Momordica charantia*) improves insulin sensitivity by increasing skeletal muscle insulin-stimulated IRS-1 tyrosine phosphorylation in high-fat-fed rats." *Br. J. Nutr.* 2008; 99(4): 806-12.

Fernandes, N., et al. "An experimental evaluation of the antidiabetic and antilipidemic properties of a standardized *Momordica charantia* fruit extract." *BMC Complement. Altern. Med.* 2007 Sep 24; 7:29.

Huang, H., et al. "Bitter melon (*Momordica charantia* L.) inhibits adipocyte hypertrophy and down regulates lipogenic gene expression in adipose tissue of diet-induced obese rats." *Br. J. Nutr.* 2008; 99(2): 230-9.

Dans, A., et al. "The effect of *Momordica charantia* capsule preparation on glycemic control in type 2 diabetes mellitus needs further studies." *J. Clin. Epidemiol.* 2007; 60(6): 554-9.

Xiang, L., et al. "The reparative effects of *Momordica charantia* Linn. extract on HIT-T15 pancreatic beta-cells." *Asia Pac. J. Clin. Nutr.* 2007; 16 Suppl 1: 249-52.

Roffey, B. et al. "Water extracts from *Momordica charantia* increase glucose uptake and adiponectin secretion in 3T3-L1 adipose cells." *J. Ethnopharmacol.* 2007 May; 112(1): 77-84.

Oishi, Y., et al. "Inhibition of increases in blood glucose and serum neutral fat by *Momordica charantia* saponin fraction." *Biosci. Biotechnol. Biochem.* 2007; 71(3): 735-40.

Omar, S., et al. "Hypoglycemic effect of the seeds of *Momordica charantia*." *Fitoterapia.* 2007; 78(1): 46-7.

Ojewole, J., et al. "Hypoglycaemic and hypotensive effects of *Momordica charantia* Linn (Cucurbitaceae) whole-plant aqueous extract in rats." *Cardiovasc. J. S. Afr.* 2006 Sep-Oct; 17(5): 227-32.

Mahomoodally, M., et al. "Effect of exogenous ATP on *Momordica charantia* Linn. (Cucurbitaceae) induced inhibition of d-glucose, l-tyrosine and fluid transport across rat everted

intestinal sacs *in vitro*." *J. Ethnopharmacol.* 2007 Mar; 110(2): 257-63.

Lans, C. "Ethnomedicines used in Trinidad and Tobago for urinary problems and Diabetes mellitus." *J. Ethnobiol. Ethnomedicine.* 2006 Oct 13; 2: 45.

Chuang, C., et al. "Fractionation and identification of 9c, 11t, 13t-conjugated linolenic acid as an activator of PPAR alpha in bitter gourd (*Momordica charantia* L.)." *J. Biomed. Sci.* 2006 Nov; 13(6): 763-72.

Krawinkel, M., et al. "Bitter gourd (*Momordica charantia*): A dietary approach to hyperglycemia." *Nutr. Rev.* 2006; 64(7 Pt 1): 331-7.

Harinantenaina, L., et al. "*Momordica charantia* constituents and antidiabetic screening of the isolated major compounds." *Chem. Pharm. Bull.* 2006; 54(7): 1017-21.

Abd El Sattar, E., et al. "Some toxicological studies of *Momordica charantia* L. on albino rats in normal and alloxan diabetic rats." *J. Ethnopharmacol.* 2006 Nov; 108(2): 236-42.

Yibchok-Anun, S., et al. "Slow acting protein extract from fruit pulp of *Momordica charantia* with insulin secretagogue and insulinomimetic activities." *Biol. Pharm. Bull.* 2006 Jun; 29(6): 1126-31.

Jung, M., et al. "Antidiabetic agents from medicinal plants." *Curr. Med. Chem.* 2006; 13(10): 1203-18.

Kumar, G., et al. "Effect of bitter gourd and spent turmeric on constituents of glycosaminoglycans in different tissues in streptozotocin induced diabetic rats." *Mol. Cell. Biochem.* 2006 Jun; 286(1-2): 53-8.

Reyes, B., et al. "Anti-diabetic potentials of *Momordica charantia* and *Andrographis paniculata* and their effects on estrous cyclicity of alloxan-induced diabetic rats." *J. Ethnopharmacol.* 2006 Apr; 105(1-2): 196-200.

Khan, B., et al. "Hypoglycemic activity of aqueous extract of some indigenous plants." *Pak. J. Pharm. Sci.* 2005; 18(1): 62-4.

Zheng, Z.X., et al. "The hypoglycemic effects of crude polysaccharides extract from *Momordica charantia* in mice." *Wei Sheng Yan Jiu.* 2005 May; 34(3): 361-3.

Sathishsekar, D., et al. "Beneficial effects of *Momordica charantia* seeds in the treatment of STZ-induced diabetes in experimental rats." *Biol. Pharm. Bull.* 2005; 28(6): 978-83.

Shetty, A. K., et al. "Effect of bitter gourd (*Momordica charantia*) on glycaemic status in streptozotocin induced diabetic rats." *Plant Foods Hum. Nutr.* 2005 Sep; 60(3): 109-12.

Kumar Shetty, A., et al. "Bitter gourd (*Momordica charantia*) modulates activities of intestinal and renal disaccharidases in streptozotocin-induced diabetic rats." *Mol. Nutr. Food Res.* 2005; 49(8): 791-6.

Chaturvedi, P., et al. "Effect of *Momordica charantia* on lipid profile and oral glucose tolerance in diabetic rats." *Phytother. Res.* 2004; 18(11): 954-6.

Vikrant, V., et al. "Treatment with extracts of *Momordica charantia* and *Eugenia jambolana* prevents hyperglycemia and hyperinsulinemia in fructose fed rats." *J. Ethnopharmacol.* 2001;

76(2): 139-43.

Miura, T., et al. "Hypoglycemic activity of the fruit of the *Momordica charantia* in type 2 diabetic mice." *J. Nutr. Sci. Vitaminol.* 2001; 47(5): 340-44.

Raza, H., et al. "Modulation of xenobiotic metabolism and oxidative stress in chronic streptozotocin-induced diabetic rats fed with *Momordica charantia* fruit extract." *J. Biochem. Mol. Toxicol.* 2000; 14(3): 131-39.

Ahmad, N., et al. "Effect of *Momordica charantia* (Karolla) extracts on fasting and postprandial serum glucose levels in NIDDM patients." *Bangladesh Med. Res. Counc. Bull.* 1999; 25(1): 11-13.

Ahmed, I., et al. "Effects of *Momordica charantia* fruit juice on islet morphology in the pancreas of the streptozotocin-diabetic rat." *Diabetes Res. Clin. Pract.* 1998; 40(3): 145-51.

Sarkar, S., et al. "Demonstration of the hypoglycemic action of *Momordica charantia* in a validated animal model of diabetes." *Pharmacol. Res.* 1996; 33(1): 1-4.

Ali, L., et al. "Studies on hypoglycemic effects of fruit pulp, seed and whole plant of *Momordica charantia* on normal and diabetic model rats." *Planta Med.* 1993; 59(5): 408-12.

Akhtar, M. S. "Trial of *Momordica charantia* Linn (Karela) powder in patients with maturity-onset diabetes." *J. Pak. Med. Assoc.* 1982; 32(4): 106-7.

Return to the Rain-Tree [Tropical Plant Database File on Bitter Melon](#)

Copyrighted 2025 by [Leslie Taylor](#). All rights reserved.