

Anti-Parasitic, Larvicidal & Anti-Malarial Actions of Pau d'arco (*Tabebuia impetiginosa*)

Published Research:

- Rigamonti, G., et al. "Selective activity of *Tabebuia avellanedae* against *Giardia duodenalis* infecting organoid-derived human gastrointestinal epithelia." *Int. J. Parasitol. Drugs Drug. Resist.* 2025 Apr; 27: 100583.
- Santos, L., et al. "Larvicidal activity of phytochemicals from *Handroanthus impetiginosus* seeds against *Culex quinquefasciatus*." *Chem. Biodivers.* 2025 Jul: e01126.
- Montagnini, D., et al. "The schistosomicidal activity of ethanolic extracts from branches, leaves, flowers and fruits of *Handroanthus impetiginosus* (Mart. ex DC.) Mattos (Bignoniaceae) plant and metabolic profile characterization by UPLC-ESI-QTOF analysis." *Braz. J. Biol.* 2023 Nov; 83: e275824.
- Ramos-Milaré, A., et al. "*In vitro* effects of lapachol and β-lapachone against *Leishmania amazonensis*." *Braz. J. Med. Biol. Res.* 2023 May; 56: e12693.
- Ramos-Milaré, Á., et al. "The anti-leishmania potential of bioactive compounds derived from naphthoquinones and their possible applications: A systematic review of animal studies." *Parasitol. Res.* 2022 May; 121(5): 1247-1280.
- Peixoto, J., et al. "Epoxy-α-lapachone (2,2-Dimethyl-3,4-dihydro-spiro[2H-naphtho[2,3-b]pyran-10,2'-oxirane]-5(10H)-one): a promising molecule to control infections caused by protozoan parasites." *Braz. J. Infect. Dis.* 2023 Mar-Apr; 27(2): 102743.
- Rani, Ret al. "Mechanism of action and implication of naphthoquinone as potent anti-trypanosomal drugs." *Curr. Top. Med. Chem.* 2022; 22(25): 2087-2105.
- Rani, R., et al. "Natural naphthoquinones and their derivatives as potential drug molecules against trypanosome parasites." *Chem. Biol. Drug Des.* 2022 Dec; 100(6): 786-817.
- Virendra, S., et al. "Natural antioxidants as additional weapons in the fight against malarial parasite." *Curr. Top. Med. Chem.* 2022; 22(24): 2045-2067.
- Peixoto, J., et al. "In silico insights into the mechanism of action of epoxy-α-lapachone and epoxymethyl-lawsone in *Leishmania spp.*" *Molecules.* 2021 Jun; 26(12): 3537.
- do Nascimento, M., et al. "In silico, *in vitro* and *in vivo* evaluation of natural Bignoniaceous naphthoquinones in comparison with atovaquone targeting the selection of potential antimalarial candidates." *Toxicol. Appl. Pharmacol.* 2020 Aug; 401: 115074.

- Mohammadi, S., et al. "Medicinal plants used in the treatment of Malaria: A key emphasis to *Artemisia*, *Cinchona*, *Cryptolepis*, and *Tabebuia* genera." *Phytother. Res.* 2020 Jul; 34(7): 1556-1569.
- Pertino, M., et al. "Synthesis, trypanocidal and anti-leishmania activity of new triazole-lapachol and nor-lapachol hybrids." *Bioorg. Chem.* 2020 Oct; 103: 104122.
- Cunha Araújo, I., et al. "Efficacy of lapachol on treatment of cutaneous and visceral leishmaniasis." *Exp. Parasitol.* 2019 Apr; 199: 67-73.
- Mata-Santos, T., "Anthelmintic activity of lapachol, β-lapachone and its derivatives against *Toxocara canis* larvae." *Rev. Inst. Med. Trop.* 2015 May-Jun; 57(3): 197-204.
- Kim, M., et al. "Larvicidal activity of the active constituent isolated from *Tabebuia avellanedae* bark and structurally related derivatives against three mosquito species." *J. Agric. Food Chem.* 2013 Nov; 61(45): 10741-5.
- González-Coloma, A., et al. "Antileishmanial, antitrypanosomal, and cytotoxic screening of ethnopharmacologically selected Peruvian plants." *Parasitol. Res.* 2012 Apr; 110(4): 1381-92.
- Silva, T., et al. "Molluscicidal activities of six species of Bignoniaceae from north-eastern Brazil, as measured against *Biomphalaria glabrata* under laboratory conditions." *Ann. Trop. Med. Parasitol.* 2007 Jun; 101(4): 359-65.
- Ferreira, V. F., et al. "Trypanocidal agents with low cytotoxicity to mammalian cell line: a comparison of the theoretical and biological features of lapachone derivatives." *Bioorg. Med. Chem.* 2006 Aug; 14(16): 5459-66.
- Silva, R. S., et al. "Synthesis of naphthofuranquinones with activity against *Trypanosoma cruzi*." *Eur. J. Med. Chem.* 2006 Apr; 41(4): 526-30.
- Menna-Barreto, R. F., et al. "Effect of a beta-lapachone-derived naphthoimidazole on *Trypanosoma cruzi*: identification of target organelles." *J. Antimicrob. Chemother.* 2005 Dec; 56(6): 1034-41.
- Perez-Sacau, E., et al. "Antiplasmodial activity of naphthoquinones related to lapachol and beta-lapachone." *Chem. Biodivers.* 2005; 2(2): 264-74.
- Lima, N. M., et al. "Antileishmanial activity of lapachol analogues." *Mem. Inst. Oswaldo Cruz.* 2004 Nov; 99(7): 757-61.
- de Andrade-Neto, V. F., et al. "Antimalarial activity of phenazines from lapachol, beta-lapachone and its derivatives against *Plasmodium falciparum* in vitro and *Plasmodium berghei* in vivo." *Bioorg. Med. Chem. Lett.* 2004 Mar; 14(5): 1145-9.
- Pinto, C. N., et al. "Chemical reactivity studies with naphthoquinones from *Tabebuia* with anti-trypanosomal efficacy." *Arzneimittelforschung.* 2000; 50(12): 1120-8.
- Austin, F. R. "*Schistosoma mansoni* chemoprophylaxis with dietary lapachol." *Am. J. Trop. Med. Hyg.* 1979; 23: 412-19.
- Gilbert, B., et al. "Schistosomiasis. Protection against infection by terpenoids." *An. Acad. Brasil. Cienc.* 1970; 2 (Suppl): 397-400.

**Return to the [Tropical Database file for Pau d'arco](#)
Copyrighted 2025 by Leslie Taylor. All rights reserved.**