

Anti-Inflammatory & Muscle Relaxant Actions of Picão Preto (*Bidens pilosa*)

Published Research:

Tsuruta, K., et al. "Anti-inflammatory effects of Miyako *Bidens pilosa* in a mouse model of amyotrophic lateral sclerosis and lipopolysaccharide-stimulated BV-2 microglia." *Int. J. Mol. Sci.* 2023 Sep; 24(18): 13698.

Yan, Z., et al. "Bioactive polyacetylenes from *Bidens pilosa* L and their anti-inflammatory activity." *Nat. Prod. Res.* 2022 Dec; 36(24): 6353-6358.

Xin, Y., et al. "Anti-inflammatory activity and mechanism of isookanin, isolated by bioassay-guided fractionation from *Bidens pilosa* L." *Molecules.* 2021 Jan; 26(2): 255.

Lee, H., et al. "Anti-neuroinflammatory and anti-inflammatory activities of phenylheptatriyne isolated from the flowers of *Coreopsis lanceolata* L. via NF- κ B inhibition and HO-1 expression in BV2 and RAW264.7 Cells." *Int. J. Mol. Sci.* 2021 Jul; 22(14): 7482.

Quaglio, A., et al. "*Bidens pilosa* (Black Jack) standardized extract ameliorates acute TNBS-induced intestinal inflammation in rats." *Planta Med.* 2020; 86: 319–330.

Ocheng, F., et al. "In vitro cytotoxicity and effects on IL-1 β -Induced proinflammatory mediators by human gingival fibroblasts." *Evid. Based Complement. Alternat. Med.* 2016; 2016: 5357689.

Shen, A., et al. "Total flavonoids of *Bidens bipinnata* L. ameliorate experimental adjuvant-induced arthritis through induction of synovial apoptosis." *BMC. Complement. Altern. Med.* 2015 Dec; 15(1): 437.

Bastos, C., et al. "Use of *Bidens pilosa* L. (Asteraceae) and *Curcuma longa* L. (Zingiberaceae) to treat intestinal mucositis in mice: Toxicopharmacological evaluations." *Toxicol. Rep.* 2015 Oct; 3: 279-287.

Fotso, A., et al. "Analgesic and antiinflammatory activities of the ethyl acetate fraction of *Bidens pilosa* (Asteraceae)." *Inflammopharmacology.* 2014 Apr; 22(2): 105-14.

Liu, J., et al. "Effect of floral sources on the antioxidant, antimicrobial, and anti-inflammatory activities of honeys in Taiwan." *Food Chem.* 2013 Aug; 139(1-4): 938-43.

Yoshida, N., et al. "*Bidens pilosa* suppresses interleukin-1 β -induced cyclooxygenase-2 expression through the inhibition of mitogen activated protein kinases phosphorylation in normal human dermal fibroblasts." *J. Dermatol.* 2006; 33(10): 676-83.

Chiang, Y., et al. "Ethyl caffeate suppresses NF- κ B activation and its downstream inflammatory mediators, iNOS, COX-2, and PGE2 in vitro or in mouse skin." *Br. J. Pharmacol.* 2005 Oct; 146(3): 352-63.

Nguelefack, T., et al. "Relaxant effects of the neutral extract of the leaves of *Bidens pilosa* Linn on isolated rat vascular smooth muscle." *Phytother. Res.* 2005; 19(3): 207-10.

Chang, C., et al. "The distinct effects of a butanol fraction of *Bidens pilosa* plant extract on the

development of Th1-mediated diabetes and Th2-mediated air way inflammation in mice." *J. Biomed. Sci.* 2005; 12(1): 79-89.

Pereira, R., et al. "Immunosuppressive and anti-inflammatory effects of methanolic extract and the polyacetylene isolated from *Bidens pilosa* L." *Immunopharmacology*. 1999; 43(1): 31-7.

Jager, A., et al. "Screening of Zulu medicinal plants for prostaglandin-synthesis inhibitors" *J. Ethnopharmacol.* 1996; 52(2): 95-100.

Chih, H., et al. "Anti-inflammatory activity of Taiwan folk medicine 'ham-hong-chho' in rats." *Am. J. Chin. Med.* 1995; 23(3-4): 273-78.

Return to the [Tropical Database file for Picão Preto](#)

Copyrighted 2025 by Leslie Taylor. All rights reserved.